



UNIVERSITI PUTRA MALAYSIA

**ECOLOGICAL CHARACTERISTICS OF
APHIDOPHAGOUS *MENOCHILUS SEXMACULATUS*
FABRICIUS AND ITS PERFORMANCE AGAINST
APHIS GOSSYPHII GLOVER**

**FRANSISCUS XAVERIUS WAGIMAN
ADISUBROTO**

FP 1996 16

ECOLOGICAL CHARACTERISTICS OF
APHIDOPHAGOUS *MENOCHILUS SEXMACULATUS*
FABRICIUS AND ITS PERFORMANCE AGAINST
APHIS GOSSYPII GLOVER

By

FRANSISCUS XAVERIUS WAGIMAN ADISUBROTO

Dissertation Submitted in Fulfilment of the Requirements for
the Degree of Doctor of Philosophy in the Faculty of
Agriculture,
Universiti Pertanian Malaysia

October, 1996



ACKNOWLEDGEMENTS

I wish to express my deep appreciation to members of my supervisory committee, Professor Dr. Mohd. Yusof Hussein (Chairman), Associate Professor Dr. Ahmad Said Sajap, Dr. Rita Muhamad and Dr. Azhar Ismail (MARDI), for their untiring and continued guidance, assistance, critical discussions and encouragement in the preparation of this dissertation.

I would like to thank Head of the Department of Plant Protection and Dean of the Faculty of Agriculture, Universiti Pertanian Malaysia (UPM), for granting permission to use facilities for carrying out laboratory and field experiments.

My special thank to members of the Entomological Laboratory, Department of Plant Protection, Faculty of Agriculture, UPM, particularly Mr. Hassanjaya Yahya, for his technical assistance during the conduct of the laboratory and field experiments.

I am indebted to the staff of the Experimental Field II Section, UPM, particularly Mr. Sahrir, for his technical assistance during the conduct of the field experiment.

The help and critical views of Associate Professor Dr. Yusof Ibrahim, Department of Plant Protection, UPM, is gratefully acknowledged.



Thanks are due to my friends and fellow graduate students, UPM, for their co-operation and discussion throughout my study period.

I am extremely grateful to the Dean of Faculty of Agriculture, Universitas Gadjah Mada (UGM), Yogyakarta, Indonesia, for his support and the Rector of UGM for granting the study leave.

I wish to thank the SEAMEO Regional Center for Graduate Study and Research in Agriculture (SEARCA) for awarding the scholarship for my study.

To my wife, Lilis, and my children - Agung, Beni, Candra, I thank them for their love and encouragement throughout my stay in Malaysia.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	xii
LIST OF FIGURES	xv
LIST OF PLATES	xviii
GLOSSARY AND ABBREVIATIONS	xix
ABSTRACT	xxi
ABSTRAK	xxiii
 CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW.....	7
Aphids Infesting Chilli in Malaysia	7
Aphid Species	7
Plant Damage	8
Vector Control	9
Aphid Predators	10
Aphidophagous <i>M. sexmaculatus</i>	11
Distribution	11
Prey Finding	13
Prey Requirement	14
Predator Fecundity	14
Criteria of a Good Aphid Predator	15



	Page
Predator-Prey Relationship	17
Functional Response	17
Numerical Response	20
Prey Preference	21
Predation Models	22
III GENERAL MATERIALS AND METHODS	27
Host Plants	27
Groundnut	27
Corn	32
Chilli	32
Sambau Weed	32
Culture of Aphids	33
Culture of <i>M. sexmaculatus</i>	33
IV LIFE HISTORY AND AGE STRUCTURE OF <i>M. SEXMACULATUS</i> AND <i>A. GOSSYPHII</i>	37
Introduction	37
Materials and Methods	38
Life History of <i>M. sexmaculatus</i> and <i>A. gossypii</i>	38
Age Structure of <i>M. sexmaculatus</i> and <i>A. gossypii</i>	39
Age Structure of <i>M. sexmaculatus</i>	41
Age Structure of <i>A. gossypii</i>	42
Statistical Analysis	43
Results	44
Life History of <i>M. sexmaculatus</i> and <i>A. gossypii</i>	44
Age Structure of <i>M. sexmaculatus</i> and <i>A. gossypii</i>	46
Age Structure of <i>M. sexmaculatus</i>	46
Age Structure of <i>A. gossypii</i>	49
Discussion	49
Conclusions	53

	Page
V SURVIVORSHIP AND FERTILITY OF <i>M. SEXMACULATUS</i> AND <i>A. GOSSYPHII</i>	54
Introduction	54
Materials and Methods	55
Life Table Construction	55
Hatchability of <i>M. sexmaculatus</i> Eggs	57
Statistical Analysis	57
Results	58
Survivorship and Fertility of <i>M. sexmaculatus</i>	58
Survivorship and Fertility of <i>A. gossypii</i>	60
Hatchability of <i>M. sexmaculatus</i> Eggs	64
Discussion	64
Conclusions.....	69
VI COLONISATION AND DISTRIBUTION PATTERNS OF <i>A. GOSSYPHII</i> AND <i>M. SEXMACULATUS</i> ON CHILLI	70
Introduction	70
Materials and Methods	71
Colonisation Pattern of <i>A. gossypii</i> and <i>M. sexmaculatus</i>	71
Spatial Distribution Patterns	71
Response of <i>M. sexmaculatus</i> to <i>A. gossypii</i>	74
Distribution	74
Aggregation Index of <i>M. sexmaculatus</i>	75
Results	75
Colonisation Pattern of <i>A. gossypii</i> and <i>M. sexmaculatus</i>	75
Spatial Distribution Patterns	77
Response of <i>M. sexmaculatus</i> to <i>A. gossypii</i>	82
Distribution	82
Aggregation of <i>M. sexmaculatus</i>	82
Discussion	85

	Page
Conclusions.....	87
VII PREY REQUIREMENT FOR <i>M. SEXMACULATUS</i>	88
Introduction	84
Materials and Methods	89
Prey Requirement for <i>M. sexmaculatus</i> Larva	89
Minimum Prey Requirement	89
Maximum Prey Requirement	90
Prey Requirement for Adult <i>M. sexmaculatus</i>	90
Minimum and Optimum Prey Requirement	90
Maximum Prey Requirement	91
Results	91
Prey Requirement for <i>M. sexmaculatus</i> Larva	92
Minimum Prey Requirement	92
Maximum Prey Requirement	94
Prey Requirement for Adult <i>M. sexmaculatus</i>	94
Minimum and Optimum Prey Requirement	94
Maximum Prey Requirement	97
Discussion	97
Conclusions.....	99
VIII FUNCTIONAL AND NUMERICAL RESPONSES OF <i>M. SEXMACULATUS</i> TO <i>A. GOSSYPHII</i>	100
Introduction	100
Materials and Methods	101
Functional Response of <i>M. sexmaculatus</i>	101
Numerical Response of <i>M. sexmaculatus</i>	103
Results	104
Functional Response of <i>M. sexmaculatus</i>	104
Numerical Response of <i>M. sexmaculatus</i>	107
Discussion	112

	Page
Conclusions	114
IX PREY PREFERENCE AND CAPTURE BY <i>M. SEXMACULATUS</i>	115
Introduction	115
Materials and Methods	115
Preference for Developmental Stages of <i>A. gossypii</i> ...	116
Preference for Aphid Species	118
Results	120
Preference for Developmental Stages of <i>A. gossypii</i> ...	120
Preference for Aphid Species	124
Preference for <i>R. maidis</i> and <i>A. gossypii</i>	124
Preference for <i>A. craccivora</i> and <i>A. gossypii</i> ..	124
Discussion	126
Conclusions.....	130
X IMPACT OF <i>M. SEXMACULATUS</i> ON SUPPRESSING <i>A. GOSSYPHII</i> POPULATION	131
Introduction	131
Materials and Methods	132
Experiment 1. Synchronisation and Voracity	132
Experiment 2. Predator Impact	135
Results	137
Experiment 1. Synchronisation and Voracity	137
Experiment 2. Predator Impact	140
Discussion	144
Conclusions.....	145
XI GENERAL DISCUSSION AND CONCLUSIONS	146

	Page
Performance of <i>M. sexmaculatus</i> Against <i>A. gossypii</i>	146
Search for Prey and Aggregation to High Prey	
Density	148
Synchronisation	151
Prey Specificity	151
Voracity and Suppression of Prey Population	152
Intrinsic Rate of Increase	154
Survival in Low Prey Density	155
Further Research	157
Conclusions	157
LITERATURE CITED	160
APPENDICES	171
Appendix A. Additional Tables	172
Appendix B. Analysis of Functional Response	194
Appendix C. Computing Procedure of van Emden's Model Using Microsoft Excel Version 5.0	196
VITA	199

LIST OF TABLES

Table	Page
1 Developmental period (days) of <i>M. sexmaculatus</i> feeding on <i>A. craccivora</i> and <i>H. setariae</i> in the laboratory and developmental period (days) of <i>A. gossypii</i> reared on chilli leaf in the laboratory and field cage	45
2 Densities and proportion of each developmental stage of <i>M. sexmaculatus</i> recorded in the chilli field within 16 to 46 DAT	48
3 Densities and proportion of each developmental stage of <i>A. gossypii</i> recorded in the chilli leaves at 28 DAT	51
4 Fertility and rate of increase of <i>M. sexmaculatus</i> maintained on <i>A. craccivora</i> and <i>H. setariae</i> in the laboratory, and <i>A. gossypii</i> maintained on chilli leaf in the laboratory and field cage	63
5 Mean density and distribution parameters for <i>A. gossypii</i> within a chilli plot	78
6 Mean density and distribution parameters for <i>M. sexmaculatus</i> within a chilli plot	79
7 Fresh body weight and third instar equivalents of different developmental stages of <i>A. gossypii</i>	92
8 Parameters of the functional response referred to as Holling's and Rogers's equation, and expected maximum consumption by <i>M. sexmaculatus</i> against <i>A. gossypii</i>	107
9 Different combinations of <i>A. gossypii</i> presented as prey to <i>M. sexmaculatus</i>	117
10 Relative size of different stages of <i>M. sexmaculatus</i> , <i>A. gossypii</i> , <i>A. craccivora</i> and <i>R. maidis</i>	121



Table		Page
11	Suggested criteria for classification of preference for two prey types by a predator	130
12	Mean numbers of survival <i>A. gossypii</i> per chilli plant at various levels of predator-prey synchronisation and predator voracity	138
13	Effect of predator number and release time of <i>M. sexmaculatus</i> on the prey suppression time (days)	139
14	Mean numbers of <i>A. gossypii</i> and <i>M. sexmaculatus</i> per chilli plant within four replications and 13 observation dates, before and after release of the predator at day 15	141
15	Mean numbers of <i>A. gossypii</i> and <i>M. sexmaculatus</i> per chilli plant within cages with and without release of <i>M. sexmaculatus</i>	142
16	Analysis of variance of transformed data based on \sqrt{x} for proportion of developmental stages of <i>M. sexmaculatus</i> recorded in the chilli field within 16 to 46 DAT	173
17	Analysis of variance of transformed data based on \sqrt{x} for proportion of developmental stages of <i>A. gossypii</i> recorded in the chilli leaves at 28 DAT	173
18	Life and fertility table for <i>M. sexmaculatus</i> feeding on <i>A. craccivora</i> in the laboratory	174
19	Life and fertility table for <i>M. sexmaculatus</i> feeding on <i>H. setariae</i> in the laboratory	175
20	Life and fertility table for <i>A. gossypii</i> maintained on the chilli leaf in the laboratory	178
21	Life and fertility table for <i>A. gossypii</i> maintained on the chilli leaf in the field cage	179

Table		Page
22	Analysis of variance of transformed data based on $\arcsin \sqrt{x}$ for incubation methods of <i>M. sexmaculatus</i> eggs in the laboratory	180
23	Mean density and distribution parameters for <i>A. gossypii</i> within a chilli plant	181
24	Mean density and distribution parameters for <i>M. sexmaculatus</i> eggs within a chilli plant	182
25	Analysis of variance of linear regression between k value of <i>A. gossypii</i> and <i>M. sexmaculatus</i> distributed in a chilli plot	183
26	Analysis of variance of linear regression between k value of <i>A. gossypii</i> and <i>M. sexmaculatus</i> distributed in a chilli plant	183
27	The constants (c and μ) of relationship $[(\beta_i)=c(\alpha_i)^\mu]$ between number proportion of <i>A. gossypii</i> (α_i) and number proportion of <i>M. sexmaculatus</i> (β_i) distributed laterally within chilli plot and vertically within chilli plant	184
28	Number of <i>A. gossypii</i> provided and eaten by larvae of <i>M. sexmaculatus</i> representing minimum prey required for developing into next developmental stages	185
29	Maximum number of <i>A. gossypii</i> eaten by larvae and adults of <i>M. sexmaculatus</i>	186
30	Functional response of first instar <i>M. sexmaculatus</i> preying on first instar <i>A. gossypii</i> in the laboratory	187
31	Functional response of fourth instar <i>M. sexmaculatus</i> preying on fourth instar <i>A. gossypii</i> in the laboratory	188
32	Functional response of female <i>M. sexmaculatus</i> preying on fourth instar <i>A. gossypii</i>	189

Table		Page
33	Functional response of first instar <i>M. sexmaculatus</i> preying on mixture stages of <i>A. gossypii</i> in the field cage	190
34	Probability of capture and differences between <i>A. gossypii</i> (various instars) presented and eaten by <i>M. sexmaculatus</i> in the laboratory	191
35	Transformed data based on \sqrt{x} transformation of proportion of prey aphid eaten by <i>M. sexmaculatus</i> and calculated T values of amongst points in unity, expected and observed proportion of prey eaten	192
36	Observed time periods (days) required by <i>M. sexmaculatus</i> larvae to eliminate <i>A. gossypii</i> numbers at various predator numbers and release times of the predator	193
37	Anova of the observed time periods (days) required by <i>M. sexmaculatus</i> larvae to eliminate <i>A. gossypii</i> numbers at various predator numbers and release times of the predator...	193

LIST OF FIGURES

Figure		Page
1	Diagrammatic representation of the chilli aphid population system, showing the main components and their interactions	5
2	A schematic diagram of the processes in rearing aphids and <i>M. sexmaculatus</i>	34
3	Pattern of age structure of <i>M. sexmaculatus</i> observed in the chilli field at 16 to 46 DAT	47
4	Pattern of age structure of <i>A. gossypii</i> colonising chilli at 28 DAT	50
5	The curves of age-specific survival and fertility rates of <i>M. sexmaculatus</i> preying on <i>A. craccivora</i> (A) and on <i>H. setariae</i> (B) in the laboratory	61
6	The curves of age-specific survival and fertility rates of <i>A. gossypii</i> maintained on chilli leaf in the laboratory (A) and in the field cage (B)	62
7	Hatchability of <i>M. sexmaculatus</i> eggs collected from the laboratory (A and B) and from chilli field (C)	65
8	Percentage of chilli plants colonised by <i>A. gossypii</i> (AG) only, both AG and <i>M. sexmaculatus</i> (MS) and MS only	76
9	Distribution of <i>A. gossypii</i> and <i>M. sexmaculatus</i> (eggs) in relation to the position of the chilli leaf	81
10	The relationship between \hat{k} values of <i>A. gossypii</i> and <i>M. sexmaculatus</i> within plot and within plant	83

Figure		Page
11	The relationship $[(\beta_i) = c(\alpha_i)^u]$ between proportion of <i>A. gossypii</i> (α_i) and proportion of <i>M. sexmaculatus</i> (β_i) per chilli plant within a plot for nine observation dates (A) and per chilli leaf within a plant for 12 sample plants (B)	84
12	Minimum and maximum number of <i>A. gossypii</i> (in TIES) required by <i>M. sexmaculatus</i>	93
13	The age of <i>M. sexmaculatus</i> larva in relation to the abundance of <i>A. gossypii</i>	95
14	Prey (<i>A. gossypii</i>) requirement for an adult <i>M. sexmaculatus</i>	96
15	Holling's Type II functional response of <i>M. sexmaculatus</i> against <i>A. gossypii</i>	105
16	Fluctuation of <i>A. gossypii</i> and <i>M. sexmaculatus</i> populations during December 1993 (A) and July 1994 (B) planting season	109
17	The time-lag relationship between the density of <i>A. gossypii</i> and the density of <i>M. sexmaculatus</i> in the second planting season of July 1994	110
18	Linear regression analyses on the numerical response of <i>M. sexmaculatus</i> against <i>A. gossypii</i> showing the two types: immediate response (A) and 3-day delayed response (B)	111
19	Proportion of <i>A. gossypii</i> numbers comprising first, second, third and fourth instar presented to and eaten by first, second, third and fourth instar and adults (females) of <i>M. sexmaculatus</i> and their differences between presented and eaten	122

Figure		Page
20	Probability of capturing first, second, third and fourth instars of <i>A. gossypii</i> by first, second, third and fourth instars and females of <i>M. sexmaculatus</i>	123
21	The result of preference experiment in which <i>R. maidis</i> and <i>A. gossypii</i> were presented together in various ratios as prey for female of <i>M. sexmaculatus</i>	125
22	The result of preference experiment in which <i>A. craccivora</i> and <i>A. gossypii</i> were presented together in various ratios as prey for female of <i>M. sexmaculatus</i>	127
23	Development of <i>M. sexmaculatus</i> population in relation to the changes of <i>A. gossypii</i> population in the field cage	143

LIST OF PLATES

Plate		Page
1	A chilli plant (variety Kulai) heavily colonised by <i>A. gossypii</i>	2
2	The aphidophagous <i>M. sexmaculatus</i>	4
3	A colony of <i>A. craccivora</i> on a leaf petiole of long bean <i>Vigna sesquipedalis</i> Fruw	28
4	A colony of corn leaf aphid <i>R. maidis</i> on the leaf sheath of sweet corn	29
5	A colony of rusty plum aphid (<i>H. setariae</i>) on the sambau weed (<i>E. indica</i>)	30
6	A hydroponics system for culturing <i>A. craccivora</i> on groundnut seedlings	31
7	A cylindrical cage is used for <i>M. sexmaculatus</i> oviposition	36
8	The experimental chilli plots	40
9	An iron frame and nylon screen (32 mesh) cage (16 x 8 x 3 m) divided into eight compartments (4 x 4 x 3 m) for studying the impact of <i>M. sexmaculatus</i> on <i>A. gossypii</i> population	136

GLOSSARY AND ABBREVIATIONS

AG	: <i>Aphis gossypii</i> Glover
ANOVA	: Analysis of variance
<i>aggregation</i>	: coming together of coccinelid or aphid into a group in plants
CRD	: completely randomised design
<i>cohort</i>	: an initial number or unit of individual insects used in the study on life table in which its development is recorded from birth until the last member is dead.
<i>distribution</i>	: the position, arrangement, or frequency of occurrence (as of the members of a group) over an area or throughout a space or unit of time
DAT	: days after transplanting
DT	: doubling time
<i>efficiency</i>	: effectiveness of the predator in affecting coexistence of the prey.
<i>fecundity</i>	: a measure of the total egg production by a female (Southwood, 1978).
<i>fertility</i>	: the number of viable eggs laid by a female (Southwood, 1978).
<i>functional response</i>	: the form of an increasing number of prey eaten per predator as prey density increases, at least up to some limiting value representing maximum prey consumption within a prescribed time.
IPM	: Integrated Pest Management
L:D	: Light:Dark period
<i>life cycle</i>	: total number of days of immature stadia; egg, larva and pupa (for MS) or nymphal instars (for AG).
<i>life table</i>	: tabulation presenting complete data on the mortality and fecundity schedule of a cohort.
<i>maximum prey requirement</i>	: maximum prey eaten per day by the predator as indicator of voracity.

<i>minimum prey requirement</i>	: least number of prey required by the predator for life maintenance.
MS	: <i>Menochilus sexmaculatus</i> Fabricius
<i>numerical response</i>	: the increase in the numbers of predators as prey numbers increase.
<i>optimum prey requirement</i>	: least number of prey required by the predator to produce eggs.
<i>performance</i>	: attribute of the predator representing its ability to find and consume prey, suppress and keep prey density at levels that will not cause economic crop damage.
<i>postreproduction period</i>	: period of time (days) between the last time to produce eggs or new progenies and die.
<i>preference</i>	: choose the better prey than another represented by the deviation of the proportion of the prey attacked from the proportion available in the environment.
<i>prereproduction period</i>	: period of time (days) between emerging from pupa and the first time laying eggs.
<i>prudent predator</i>	: a character of predator to obtain its living needs accurately and judiciously
<i>reproduction period</i>	: period of time (days) to produce eggs or new progenies.
RH	: relative humidity
TIES	: third instar equivalents
T	: generation time
UPM	: Universiti Pertanian Malaysia

Abstract of dissertation submitted to the Senate of Universiti Pertanian Malaysia in fulfilment of the requirements for the degree of Doctor of Philosophy.

**ECOLOGICAL CHARACTERISTICS OF APHIDOPHAGOUS
MENOCHILUS SEXMACULATUS FABRICIUS AND ITS
PERFORMANCE AGAINST *APHIS GOSSYPHII* GLOVER**

By

FRANSISCUS XAVERIUS WAGIMAN ADISUBROTO

October, 1996

Chairman: Professor Mohd. Yusof Hussein, Ph.D.

Faculty : Agriculture

The chilli aphid, *Aphis gossypii* Glover (Homoptera: Aphididae) (here after abbreviated as AG), is vector of the chilli veinal mottle virus (CVMV) and is a main pest of chilli in Malaysia. The aphidophagous *Menochilus sexmaculatus* Fabricius (Coleoptera: Coccinellidae) (here after abbreviated as MS) is a well known predator of the aphid and hereby its ecological characteristics and predatory performance were assessed.

In nature, MS maintains its population on aphids of corn, beans, sambau weed and other plants. The predator searched its prey in random manner and tended to aggregate in high prey density. Both prey and predator were distributed

in a clumped pattern. Larval and adult MS showed no preference for any prey stages and prey species. However, there was a weak preference for AG over *Aphis craccivora* Koch.

MS was very voracious; the larva (1st, 2nd, 3rd and 4th instar) and adult consumed 8, 43, 46, 125 and 275 TIES of AG per day, respectively. The minimum prey requirement for oviposition was 50 TIES per day. The adult survived well when supplied with 10 TIES per day. The average fecundity was 378 to 745 eggs/♀, depending on aphid prey species (*A. craccivora* and *Hysteroneura setariae* (Thos.)). The predator numbers increased at the rate (r_m) of 0.1663 to 0.2116, while a higher r_m for the prey, AG, was calculated to be 0.4097 to 0.4645.

The predator exhibited a delayed density-dependent relationship with the prey as indicated by a strong functional response of Holling's Type II and a direct numerical response to AG. In the field cages, MS was able to eliminate its prey (AG) population within nine days given the initial ratio (mated female MS : AG) of 1:6,400. In the open field, AG population was suppressed completely within three weeks given the ratio (predator : prey) of 1:50. Results of the study may contribute in developing biological control programme of aphid vector in Malaysia and other countries where the MS is found.

Abstrak disertasi yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi keperluan untuk mendapatkan ijazah Doktor Falsafah.

**CIRI-CIRI EKOLOGI *MENOCHILUS SEXMACULATUS* FABRICIUS
AFIDOFAGUS DAN PRESTASINYA TERHADAP
APHIS GOSSYPHII GLOVER**

Oleh

FRANSISCUS XAVERIUS WAGIMAN ADISUBROTO

Oktober, 1996

Pengerusi: Professor Mohd. Yusof Hussein, Ph.D.

Fakulti : Pertanian

Afid cili, *Aphis gossypii* Glover (Homoptera: Aphididae) (selanjutnya disingkat sebagai AG) adalah vektor kepada virus cili yang dinamakan dengan CVMV dan merupakan perosak utama tanaman cili di Malaysia. Serangga afidofagus, *Menochilus sexmaculatus* Fabricius (Coleoptera: Coccinellidae) (selanjutnya disingkat sebagai MS) amat terkenal sebagai pemangsa afid tersebut dan di sini ciri-ciri ekologi serta prestasi pemangsaanya telah dinilai.

Dalam alam semula jadi, populasi MS bergantung hidup pada afid jagung, kacang, rumpai sambau dan tumbuh-tumbuhan lain. MS mencari mangsa secara rawak dan cenderung berkumpul pada populasi mangsa yang tinggi. Kedua-dua mangsa dan pemangsa tersebar secara berkelompok. Larva dan dewasa MS tidak memilih mana-mana peringkat mangsa dan species afid. Walau

bagaimanapun AG lebih digemari daripada *Rhopalosiphum maidis* (Fitch.) dan *Aphis craccivora* Koch.

MS didapati sangat rakus; larva (instar 1, 2, 3 dan 4) dan peringkat dewasa memakan AG sebanyak 8, 43, 46, 125 dan 275 TIES (setara instar 3) setiap hari. Untuk bertelur MS memerlukan jumlah minimum mangsa sebanyak 50 TIES setiap hari. MS dewasa hidup selesa dengan memakan 10 TIES per hari. Min kesuburan MS adalah antara 378 hingga 745 biji telur per betina, bergantung kepada species afid yang dimakan (*A. craccivora* dan *Hysteroneura setariae* (Thos.)). Bilangan pemangsa meningkat dengan kadar pertumbuhan (r_m) 0.1663 hingga 0.2116, manakala kadar pertumbuhan yang lebih tinggi untuk mangsa, AG, telah dianggarkan setinggi 0.4097 hingga 0.4645.

MS telah menunjukkan hubungan kepadatan tertakluk terlewat dengan mangsanya berdasarkan kepada respon fungsian Holling jenis II yang kuat dan respon numerik langsung terhadap AG. Di dalam sangkar, MS mampu mengawal populasi AG dalam masa sembilan hari pada nisbah awalan (MS betina yang sudah mengawan : AG) 1:6,400. Di lapangan, populasi AG dikawal sepenuhnya oleh MS dalam masa tiga minggu pada nisbah (pemangsa : mangsa) 1:50. Hasil penyelidikan ini boleh memberi sumbangan kepada perkembangan program kawalan biologi terhadap vektor afid di Malaysia dan di negara lain dimana MS didapati.

CHAPTER I

INTRODUCTION

Aphids are among the most damaging insect pests of vegetable crops. They cause economic crop loss by directly sucking sap from the plants and indirectly transmitting plant viruses. An aphid species, *Aphis gossypii* Glover (Homoptera: Aphididae) (hereafter referred to as AG; Plate 1), is a vector of more than 50 plant viruses (Blackman and Eastop, 1989). AG efficiently transmits non-persistent chilli veinal mottle virus (CVMV) (Ong et al., 1978; Lee et al., 1994) in the nursery and in the field. CVMV is among the most important constraints of chilli production in Malaysia (Syed and Loke, 1995) and was known to significantly reduce the chilli yield. The yield reduction may reach 60% if plants were infected at the early stage of growth (Ong et al., 1980).

The viral disease of crop plants may be controlled indirectly by controlling the vector. Under natural situation the vector populations are normally controlled by the natural enemies especially by polyphagous predators such as the ladybird beetle. The important role of the ladybird beetle as predators of aphids is well known since the mid-19th century (Whitcomb, 1981). Among the ladybird beetles *Menochilus sexmaculatus* Fabricius (Coleoptera: Coccinellidae) (hereafter